



BACTERIA

“If you don't like bacteria, you're on the wrong planet.”

COURSE : B.Sc BOTANY
SEMESTER : I
PAPER : MICROLOGY AND
PHYCOLOGY/ BOT CC 101
TOPIC : BACTERIA
FACULTY : HENA NAZ
EMAIL ID : henanaz64@gmail.com

Bacteria – the benign, the bad and the beautiful.

Facts :

- **Bacteria are present almost everywhere from deep in the earth's crust to the polar ice caps and oceans to inside the bodies of plants and animals.**
- **There are more bacteria in your mouth than there are people in the world.**
- **Babies are born with no bacteria in their bodies.**
- **Electronics, cellphones, laptops, keyboards etc. hold a lot of bacteria. Single smartphone screens hold 18 times more bacteria than a toilet handle.**
- **Smell of rain is caused by a bacteria called actinomycetes.**
- **Sweat itself is odourless. It's the bacteria on the skin that mingles with it and produces body odour.**
- **Horseshoe crab blood is worth US \$15000/ L due to its ability to detect bacteria.**
- **Gonorrhea bacteria is the strongest creatures on the earth as they can pull 100,000 times their own weight.**

Introduction

- Bacteria were discovered by Leeuwenhoek in 1676.
- They are the primitive forms of life.
- They are monerans and comprises a group of prokaryotic organisms which is characterized by:
 - Peptidoglycan wall
 - Compacted but naked DNA with attached mesosome
 - Reserve food material made up of glycogen and fats
 - Gas vacuoles may occur
 - All membranes bound cell organelles completely absent
 - 70S Ribosome occurs
- They have varied forms of nutrition.



**Antonie van Leeuwenhoek
(1632-1723)**

Bacterial cell structure & Cell morphology

Cocci


coccus





diplococci



diplococci
encapsulated
Pneumococcus




Staphylococci



streptococci



sarcina



tetrad

Others




enlarged rod
Fusobacterium



Vibrio



Comma's form
Bdellovibrio



Club Rod
Corynebacteriaceae



Helical form
Helicobacter pylori



Corkscrew's form
Borrelia burgdorferi



Filamentous



spirochete

Bacilli



coccobacillus



bacillus



diplobacilli



palisades



Streptobacilli

Budding and appendaged bacteria




hypha




stalk

Structure of a typical bacterial cell



Cell Envelope

- ❑ Outer covering of protoplast of bacterial cell.
- ❑ The cell envelope of bacteria consist 3 components:
 - Glycocalyx
 - cell wall
 - plasma membrane



Glycocalyx

- ☐ Some bacteria have an additional layer outside of the cell wall called the Glycocalyx.
- ☐ This outermost mucilaginous layer consist of non cellulosic polysaccharides with or without proteins.
- ☐ It occurs as
 - Slime layer when glycocalyx occur in the form of loose mucilage sheath
 - Capsule when thick and tough mucilage covering.

- ☐ Glycocalyx gives sticky character to the cell.
- ☐ It is not absolutely essential for survival of bacteria. However it has several secondary functions:
 - Prevention of desiccation
 - Protection from phagocytes
 - Protection from toxic chemicals and drugs
 - Protection from viruses
 - Attachment
 - Immunogenicity
 - Virulence

Cell wall

- ❑ The bacterial cell wall differs from that of all other organisms by the presence of peptidoglycan which is located immediately outside of the cytoplasmic membrane.
- ❑ Peptidoglycan is made up of a polysaccharide backbone consisting of alternating N-Acetylmuramic acid (NAM) and N-acetylglucosamine (NAG) residues in equal amounts.
- ❑ Peptidoglycan is responsible for the rigidity of the bacterial cell wall and for the determination of cell shape.
- ❑ It is relatively porous and is not considered to be a permeability barrier for small substrates.




Fig: The structure of peptidoglycan.

Types of bacterial cell wall

- ❑ There are two main types of bacterial cell walls, those of gram-positive bacteria and those of gram-negative bacteria, which are differentiated by their Gram staining characteristics.
- ❑ Gram-positive cell walls are thick and Gram-negative cell walls are thin peptidoglycan layer adjacent to the cytoplasmic membrane.
- ❑ Peptidoglycan layer constitutes almost 95% of the cell wall in some gram-positive bacteria and as little as 5-10% of the cell wall in gram-negative bacteria.




Fig: Structure of bacterial cell wall

Cell membrane

- ❑ The plasma membrane or bacterial cytoplasmic membrane is composed of a phospholipid bilayer and thus has all of the general functions of a cell membrane.
- ❑ It is selectively permeable.
- ❑ It is metabolically active as it takes part in respiration, synthesis of lipids and cell wall components.




Fig: Bacterial cell membrane

Mesosome

- ❑ Multi-laminated structure formed invaginations of plasma membrane.
- ❑ Principal sites of respiratory enzymes.
- ❑ Coordinate nuclear & cytoplasmic division during binary fission.
- ❑ More prominent in Gram +ve bacteria.




Fig : structure of mesosome

Ribosome

- ❑ **Bacteria** and archaeobacteria have smaller **ribosomes**, termed **70S ribosomes**, which are composed of a small 30S subunit and large 50S subunit.
- ❑ The "S" stands for svedbergs, a unit used to measure how fast molecules move in a centrifuge



Fig : structure of 70S ribosome

Nucleoid (Bacterial Chromosome)

- Since the bacterial cell is prokaryotic, a true nucleus is absent.
- The nuclear material is represented by DNA which is not associated with histones.
- The bacterial DNA is circular and is attached at a point to the plasma membrane.





Fig : nucleoid organisation in bacteria

Plasmid

- ☐ Extra nuclear genetic elements consisting of DNA.
- ☐ Transmitted to daughter cells during binary fission.
- ☐ May be transferred from one bacterium to another.
- ☐ Not essential for life of the cell.
- ☐ Confer certain properties e.g. drug resistance, toxicity.



Flagella

- ❑ Long (3to12 μ), filamentous surface appendages.
- ❑ Organs of locomotion.
- ❑ Chemically, composed of proteins called flagellins.
- ❑ The number and distribution of flagella on the bacterial surface are characteristic for a given species- hence are useful in identifying and classifying bacteria.
- ❑ Flagella may serve as antigenic determinants (e.g. the H antigens of Gram-negative enteric bacteria).
- ❑ Presence shown by motility e.g. hanging drop preparation.





Fig : structure of flagella

Types of flagellar arrangement




Polar/ Monotrichous – single flagellum at one pole



Lophotrichous – tuft of flagella at one pole



Amphitrichous – flagella at both poles





Peritrichous – flagella all over



Amphilophotrichous – tuft of flagella at both ends

Pilli and fimbriae

- ☐ Proteins filaments.
- ☐ Shorter and thinner: fimbriae
- ☐ Longer and fewer: pili
- ☐ Sex Pili : The filaments that are best known for their function in conjugation.



Inclusion bodies

- ❑ Are reserve deposits of bacterial.
- ❑ Cells accumulate nutrient when they are plenty and use when deficient.
- ❑ Major Inclusion bodies are:
 - Gas vacuoles-allows for buoyancy .
 - Sulfur granules.
 - Polyphosphate granules.
 - Glycogen granules.
 - Lipid inclusion.
 - Magnetosomes.





Thank you